


NOVAC

Network for Observation of Volcanic and Atmospheric Change

B. Galle¹, U. Platt², M. Van Roozendael³, C. Oppenheimer⁴, T. Hansteen⁵, G. Boudon⁶, M. Burton⁷, H. Delgado⁸, W. Strauch⁹, E. Malavassi¹⁰, G. Garzon¹¹, C. Pullinger¹², M. Kasereka¹³

1. Chalmers University of Technology, SE-412 96 Gothenburg, Sweden
 2. Heidelberg University, Germany
 3. Belgian Institute for Space Aeronomy, Belgium
 4. Cambridge University, United Kingdom
 5. IFM-GEOMAR Research Center, Germany
 6. Institut de Physique de Globe du Paris, France
 7. Istituto Nazionale di Geofisica e Vulcanologia, Italy
 8. Universidad Nacional Autonoma de Mexico, Mexico
 9. Instituto Nicaraguense de Estudios Territoriales, Nicaragua
 10. Observatorio Volcanologico y Sismologico de Costa Rica
 11. Instituto Colombiano de Geologia y Minería, Colombia
 12. Servicio Nacional de Estudios Territoriales, El Salvador
 13. Observatoire Volcanologique de Goma, D.R. Congo
 14. Massachusetts Institute of Technology, USA
 15. University of Maryland, Baltimore County, USA
- bo.galle@riss.chalmers.se



San Cristobal volcano, Nicaragua.




Figure 1. Schematic view of the optical layout of the Dual-Beam scanning mini-DOAS instrument. The prism and the protective cover is rotated around the optical axis of the telescope, thereby scanning the field-of-view of the instrument in a plane perpendicular to the optical axis.




Figure 2. Time-resolved measurement of emissions from San Cristobal 23 November 2002, calculated using plume height and wind speed from Dual Beam mini-DOAS measurements.




Figure 3. Measurement of plume speed during the field campaign at Mt. Etna in May 2005, using a dual-beam mini DOAS system.




Figure 4. Two-dimensional vertical concentration measurement performed on the plume from Mt Etna on 18 September 2004. The altitude in the figure is relative to the scanning systems, which were situated at 950 m above sea level. From the plot one can conclude that the centre of the plume is at around 4000 meters height above sea level. The concentration is presented in mg/m³.

Introduction

NOVAC is a recently started project, funded by European Union, with the aim to establish a global network of stations for the quantitative measurement of volcanic gas emissions. The network is based on a novel type of instrument, the Scanning Dual-beam mini-DOAS, developed within the EU-project DORSIVA. Primarily the instruments will be used to provide new parameters in the toolbox of the observatories for risk assessment, gas emission estimates and geophysical research on the local scale. In addition to this, data are exploited for other scientific purposes than local volcanic gas emissions, e.g. global estimates of volcanic gas emissions, large scale volcanic correlations, studies of climate change, studies of stratospheric ozone depletion. In particular large scale validation of satellite instruments for observing volcanic gas emissions will be possible for the first time, allowing to bring observation of volcanic gas emissions from space a significant step forward.

The Scanning Dual-beam Mini-DOAS instrument represents a major breakthrough in volcanic gas monitoring; it is capable of real-time automatic, unattended measurement of the total emission fluxes of SO₂ and BrO from a volcano with better than 5 minutes time resolution during daylight. The high time-resolution of the data enables correlations with other geophysical data, e.g. seismic data, thus significantly extending the information available for real-time risk assessment and research at the volcano. By comparing high time resolution gas emission data with emissions from neighbouring volcanoes on different geographical scales, or with other geophysical events (earthquakes, tidal waves) mechanisms of volcanic forcing may be revealed.

The spectra recorded by the instrument will also be used to derive data that complement global observation systems related to climate change and stratospheric ozone depletion research.

The Scanning Dual-beam Mini-DOAS instrument represents a major breakthrough in volcanic gas monitoring; it is capable of real-time automatic, unattended measurement of the total emission fluxes of SO₂ and BrO from a volcano with better than 5 minutes time resolution during daylight. The high time-resolution of the data enables correlations with other geophysical data, e.g. seismic data, thus significantly extending the information available for real-time risk assessment and research at the volcano.

The Scanning Dual-beam Mini-DOAS instrument represents a major breakthrough in volcanic gas monitoring; it is capable of real-time automatic, unattended measurement of the total emission fluxes of SO₂ and BrO from a volcano with better than 5 minutes time resolution during daylight. The high time-resolution of the data enables correlations with other geophysical data, e.g. seismic data, thus significantly extending the information available for real-time risk assessment and research at the volcano.

The Scanning Dual-beam Mini-DOAS instrument represents a major breakthrough in volcanic gas monitoring; it is capable of real-time automatic, unattended measurement of the total emission fluxes of SO₂ and BrO from a volcano with better than 5 minutes time resolution during daylight. The high time-resolution of the data enables correlations with other geophysical data, e.g. seismic data, thus significantly extending the information available for real-time risk assessment and research at the volcano.

The Scanning Dual-beam Mini-DOAS instrument represents a major breakthrough in volcanic gas monitoring; it is capable of real-time automatic, unattended measurement of the total emission fluxes of SO₂ and BrO from a volcano with better than 5 minutes time resolution during daylight. The high time-resolution of the data enables correlations with other geophysical data, e.g. seismic data, thus significantly extending the information available for real-time risk assessment and research at the volcano.

The Scanning Dual-beam Mini-DOAS instrument represents a major breakthrough in volcanic gas monitoring; it is capable of real-time automatic, unattended measurement of the total emission fluxes of SO₂ and BrO from a volcano with better than 5 minutes time resolution during daylight. The high time-resolution of the data enables correlations with other geophysical data, e.g. seismic data, thus significantly extending the information available for real-time risk assessment and research at the volcano.

The Scanning Dual-beam Mini-DOAS instrument represents a major breakthrough in volcanic gas monitoring; it is capable of real-time automatic, unattended measurement of the total emission fluxes of SO₂ and BrO from a volcano with better than 5 minutes time resolution during daylight. The high time-resolution of the data enables correlations with other geophysical data, e.g. seismic data, thus significantly extending the information available for real-time risk assessment and research at the volcano.

The Scanning Dual-beam Mini-DOAS instrument represents a major breakthrough in volcanic gas monitoring; it is capable of real-time automatic, unattended measurement of the total emission fluxes of SO₂ and BrO from a volcano with better than 5 minutes time resolution during daylight. The high time-resolution of the data enables correlations with other geophysical data, e.g. seismic data, thus significantly extending the information available for real-time risk assessment and research at the volcano.

The Scanning Dual-beam Mini-DOAS instrument represents a major breakthrough in volcanic gas monitoring; it is capable of real-time automatic, unattended measurement of the total emission fluxes of SO₂ and BrO from a volcano with better than 5 minutes time resolution during daylight. The high time-resolution of the data enables correlations with other geophysical data, e.g. seismic data, thus significantly extending the information available for real-time risk assessment and research at the volcano.

The Scanning Dual-beam Mini-DOAS instrument represents a major breakthrough in volcanic gas monitoring; it is capable of real-time automatic, unattended measurement of the total emission fluxes of SO₂ and BrO from a volcano with better than 5 minutes time resolution during daylight. The high time-resolution of the data enables correlations with other geophysical data, e.g. seismic data, thus significantly extending the information available for real-time risk assessment and research at the volcano.

The Scanning Dual-beam Mini-DOAS instrument represents a major breakthrough in volcanic gas monitoring; it is capable of real-time automatic, unattended measurement of the total emission fluxes of SO₂ and BrO from a volcano with better than 5 minutes time resolution during daylight. The high time-resolution of the data enables correlations with other geophysical data, e.g. seismic data, thus significantly extending the information available for real-time risk assessment and research at the volcano.

The Scanning Dual-beam Mini-DOAS instrument represents a major breakthrough in volcanic gas monitoring; it is capable of real-time automatic, unattended measurement of the total emission fluxes of SO₂ and BrO from a volcano with better than 5 minutes time resolution during daylight. The high time-resolution of the data enables correlations with other geophysical data, e.g. seismic data, thus significantly extending the information available for real-time risk assessment and research at the volcano.




Figure 5. The relationship between SO₂ and BrO measured on 5 different volcanoes measured with a scanning mini-DOAS instrument [Bobrowski et al 2003].




Figure 6. Map showing the 15 volcanoes involved in NOVAC (circles). Also shown are the current coverage of the Network for the Detection of Stratospheric Change. Note that very few NDSC stations are located near the equator.




Figure 7. Examples of SO₂ plumes of various volcanoes measured by satellite during the period 1996-2002. Note that 4 of these volcanoes are part of the NOVAC network. [Khokar et al 2004]

Institute	Volcanoes
Institut de Physique de Globe de Paris	Piton de Fournaise La Soufrière, Guadeloupe
Istituto Nazionale di Geofisica e Vulcanologia, Italy	Mount Etna
Universidad Nacional Autonoma de Mexico	Popocatepetl Fuego de Colima
Instituto Nicaraguense de Estudios Territoriales	San Cristobal Masaya
Observatorio Volcanologico y Sismologico de Costa Rica	Arenal Poas
Instituto Colombiano de Geologia y Minería	Galeras Nevado del Ruiz
Servicio Nacional de Estudios Territoriales, El Salvador	Santa Ana San Miguel
Goma Volcano Observatory, Dem Rep. Congo	Niyirangongo Niyamuragira

Table 1. Institutes and volcanoes presently constituting the NOVAC network.



This project is funded by The European Union, FP6, Global Change and Ecosystems, Natural Disasters program.